CELL

APPLICATIONS, INC.

Published on Cell Applications (https://www.cellapplications.com)

Home > Anti-Ret: Polyclonal Ret Antibody

Anti-Ret: Polyclonal Ret Antibody

- Description
- Details
- Products
- Resources

Product Sheet CC1033

Description

BACKGROUND RET (Rearranged during Transfection) encodes a receptor tyrosine kinase with key roles in cell growth, differentiation, and survival. It consists of three functional domains, an extracellular domain with four cadherin-like repeats, a cysteine-rich region, and an intracellular tyrosine kinase region . Following ligand binding to the extracellular region, receptor dimerization is mediated by activation of the cysteine-rich region that lies adjacent to the plasma membrane. Receptor dimerization leads to autophosphorylation of intracellular tyrosine residues, which subsequently activate downstream pathways of signal transduction.¹ Ligands for RET include those in the glial cell-line derived neurotrophic factor (GDNF) family, including persepherin, artemin, and neurturin. RET signals through multiple downstream pathways. One such pathway, the RAS/MEK/ERK pathway promotes cell cycle progression; another downstream pathway, P13K/AKT/NF-?B, leads to increased cell motility, survival, and progression through the cell cycle.² RET activation also stimulates p38, MAPK, JAK/STAT, and protein kinase C.³ RET is expressed widely in neural-crest derived tissues such as noradrenergic and dopamanergic neurons and neuroendocrine tissues including thyroid C cells and the adrenal medulla. RET is also involved in the development of the enteric nervous system and the kidney.

Activating RET germline mutations have been identified as the primary cause of the hereditary medullary thyroid carcinoma.3 The RET proto-oncogene has become the rational target for molecularly designed drug therapy.

REFERENCES

- 1. Mak, Y.F. & Ponder, B.A.J.: Curr. Opin. Genet. Develop. 6:82-86, 1996.
- 2. Arighi, E. et al.: Cytokine Growth Factor Rev. 16: 441-67, 2005.
- 3. Santoro, M. et al.: Eur. J Endocrinol. 133: 513-522, 1995.

Products are for research use only. They are not intended for human, animal, or diagnostic applications.



(Click to Enlarge) **Top:** TT cell lysate was subjected to Western Blot analysis using Ret antibody. **Bottom:** This antibody stains paraffin-embedded human thyroid tumor tissue in immunohistochemical analysis

Details

Cat.No.:	CC1033		
Antigen:	sequence near the human Ret carboxyl terminal.		
Isotype:	Affinity purified rabbit IgG		
Species & predicted species cross- reactivity ():	Human		
Applications & Suggested starting dilutions:	WB 1:1000 IP n/d IHC (Paraffin) n/d ICC n/d FACS n/d		
Predicted Molecular Weight of protein:	175 kDa		
Specificity/Sensitivity:	This antibody detects human Ret proteins in transfected cell lysate.		
Storage:	Store at 4° C for frequent use; at -20° C for at least one year.		

*Optimal working dilutions must be determined by end user.

Products

Resources/Documents

Product Sheet CC1033 [2]

Misc. Links

•	Site
•	Privacy
•	Returns
•	Shipping
•	Terms
•	Disclaimer
•	Distributors

Contact Us

Cell Applications, Inc

6455 Weathers Place San Diego, CA 92121 Open M-F, 8am-5pm PST

800-645-0848 info@cellapplications.com

Socialize With Us

Newsletter Signup

Subscribe to our newsletter

Source URL:https://www.cellapplications.com/anti-ret-polyclonal-ret-antibody

Links

•

[1] https://www.cellapplications.com/sites/default/files/images_product_type/FileCat000iklftci.gif
[2] https://www.cellapplications.com/sites/default/files/documents/product-sheets/Product Sheet CC1033
Ret.pdf